



Nutrients PLUS® Fertilizer with 0.37% Prodiamine

16-2-3

For pre-emergence control of grass and broadleaf weeds in lawns and ornamental turf (excluding golf course putting greens)

Landscape ornamentals, established perennials and wildflower plantings

Stops many weed grasses before they grow - Prevents crabgrass, annual bluegrass, foxtail

ACTIVE INGREDIENT:
Prodiamine: (CAS # 29091-21-2) 0.37%

OTHER INGREDIENTS: 99.63%

TOTAL: 100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See back panel for further precautions and First Aid Information below

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or repeated skin contact with product may cause allergic reactions in some individuals.

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> Call a poison control center or physician immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or physician. Do not give anything by mouth to an unconscious person.
IF INHALED:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact The National Pesticide Telecommunications Network at 1-800-858-7378 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

GUARANTEED ANALYSIS	
Total Nitrogen (N)	16.00%
2.50% Ammoniacal Nitrogen	
9.5% Urea Nitrogen†	
1.60% Other Water Soluble Nitrogen	
2.40% Water Insoluble Nitrogen*	
Available Phosphate (P ₂ O ₅)	2.00%
Soluble Potash (K ₂ O)	3.00%
Calcium (Ca)	2.00%
Sulfur (S)	3.00%
3.00% Combined Sulfur (S)	
Iron (Fe)	1.00%
Derived from poultry manure, biosolids, ammonium sulfate, urea, sulfur coated urea, muriate of potash.	
* 2.40% slowly available water insoluble nitrogen from poultry manure and biosolids.	
† 6.00% slowly available urea nitrogen from sulfur coated urea.	
F1519	
Information concerning the raw materials composing this product can be obtained by writing to the Guarantor listed on this label, and referring to the batch number found on this container.	
Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.htm	

DIRECTION FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product is a pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in established lawns and ornamental turf (excluding golf course putting greens), landscape ornamentals and established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seeds, germination and root development. When used according to directions, it will control; barnyardgrass; bluegrass; annual (*Poa annua*); carpetweed; chickweed, common; chickweed, mouseear (from seed); crabgrass (large, smooth); crowfootgrass; cupgrass, woolly; foxtails, annual; goosegrass¹; henbit; itchgrass; johnsongrass (from seed); junglerice; knotweed; kochia; lambsquarter, common; lovegrass; panicum (Texas, fall, browntop); pigweed; purslane, common; pusley, Florida; rescuegrass³, shepherd's purse²; signalgrass, broadleaf; speedwell, Persian; sprangletop; spurge, prostrate; witchgrass; woodsorrel, yellow (from seed).

¹ In many areas a single application of 4.1 to 9.3 pounds/1,000 square feet (175 to 405 pounds/acre) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 4.1 to 6.2 pounds/1,000 square feet (175 to 270 pounds/acre), followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate (see "Minimum and Maximum Application Rate by Turfgrass Species" table in the Rates of Application section of this label).

² For this weed, apply in late summer, fall or winter prior to germination.


³ Suppression only.

NET WT 50 lb (22.7 kg)

EPA Reg. No. 53883-172-82898
EPA Est. No. 047956-PA-001(A); 072744-OK-001(B)

Distributed & Guaranteed by Nutrients PLUS®, LLC
2133 Upton Dr., Ste. 126-509 • Virginia Beach, VA 23454 • 888-675-7515

WHEN TO APPLY TO ESTABLISHED TURF:
This product must be applied prior to germination of weed seeds, as it will not control established weeds.



APPLICATION DIRECTIONS:
Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days of application.

USE PRECAUTIONS:
The following precautions apply to the use of this product in turf grasses and lawns. Application of this product may thin emerged annual bluegrass and newly overseeded grasses. Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil).

Do not apply newly set sod until the following year. Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. Disturbing the herbicide barrier with cultural practices (such as core aeration) may result in reduced weed control. Do not apply to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species. Do not apply through any type of irrigation system. Do not apply aerially. Do not graze or feed livestock forage cut from treated areas.

RATES OF APPLICATION:
This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not exceed the maximum yearly rate listed:

MINIMUM AND MAXIMUM APPLICATION RATES (by Turf Grass Species)		
Turf Species	Minimum Rate	Maximum Annual Rate ¹
Bahiagrass, Bermudagrass ² Centipedegrass, Zoysiagrass, Seashore Paspalum, St. Augustinegrass Tall Fescue (including turf-Type)	2.7 lbs/1,000 sq. ft. (118 lbs/acre)	9.3 lbs/1,000 sq. ft. per year (405 lbs/acre per year)
Buffalograss Kentucky Bluegrass Perennial Ryegrass	2.7 lbs/1,000 sq. ft. (118 lbs/acre)	6.2 lbs/1,000 sq. ft. per year (270 lbs/acre per year)
Creeping Red Fescue	2.7 lbs/1,000 sq. ft. (118 lbs/acre)	4.6 lbs/1,000 sq. ft. per year (202 lbs/acre per year)
Creeping Bentgrass	2.7 lbs/1,000 sq. ft. (118 lbs/acre)	4.0 lbs/1,000 sq. ft. per year (175 lbs/acre per year)

¹ These are the maximum rates per calendar year.

² May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 3.3 pounds/1,000 square feet (145 pounds/acre). Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sprigging situations.

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of weed control.

DIRECTION FOR USE (continued)

WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit germination of turf seed if overseeded too soon after application. Follow rates and intervals in table for best overseeding/reseeding results. (See map)

USE RATE	MONTHS BEFORE OVERSEEDING		
	North	Transition	South
3 pounds/1,000 sq. ft. (135 pounds/acre)	4	4	4
4 pounds/1,000 sq. ft. (175 pounds/acre)	5	4	4
4.6 pounds/1,000 sq. ft. (202 pounds/acre)	6	5	4
4.9 pounds/1,000 sq. ft. (215 pounds/acre)		6	6
6.2 pounds/1,000 sq. ft. (270 pounds/acre)		7	7
7 pounds/1,000 sq. ft. (308 pounds/acre)			9
8 pounds/1,000 sq. ft. (350 pounds/acre)			10
9.3 pounds/1,000 sq. ft. (405 pounds/acre)			12

LANDSCAPE ORNAMENTALS (including established perennials and wildflower plantings):

This product may be applied for residual pre-emergence weed control in ornamentals. Apply at the rate of 2.7 to 9.3 pounds/1,000 square feet (118 to 405 pounds/acre) in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 9.3 pounds/1,000 square feet (405 pounds/acre) per year.

SPREADER SETTINGS to apply at the listed pounds per 1,000 square feet						
	3 lbs.	4 lbs.	4.6 lbs.	6.2 lbs.	8 lbs.	9.3 lbs.
Lely Broadcast	2.6	3.4	3.5	3.8	3.4 x 2	8.5

APPLICATION TIMING AND INFORMATION

This product may be applied to newly transplanted and established ornamentals as a broadcast, over the top or directed application. Around new transplants, delay application to allow soil to settle and water thoroughly before applying. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union. This product is a pre-emergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when applied to soil free of clods, weeds and debris such as leaves. Prior to application, control existing vegetation by hand weeding, cultivation or the use of an appropriate post-emergence herbicide.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a dry place. Store in locked area out of reach of children and domestic animals.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spilled bait, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance.

CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this container. Empty container through use according to label instructions. Do not rinse unless required for recycling. Offer for recycling, if available. If recycling is not available; then dispose of container in a sanitary landfill or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONDITIONS OF SALE AND WARRANTY

The DIRECTIONS FOR USE of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop, injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of Nutrients PLUS, LLC or the Seller. All such risks shall be assumed by the Buyer.

Nutrients PLUS, LLC warrants that this product conforms to the chemical description on the label, and is reasonably fit for the purposes set forth in the DIRECTIONS FOR USE, when it is used in accordance with such directions, subject to the inherent risks mentioned above.

Nutrients PLUS, LLC NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESSED OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. THIS WARRANTY EXTENDS TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNING, OR CAUTIONS. BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. Nutrients PLUS, LLC AND THE SELLER OFFER THIS PRODUCT, AND THE BUYER AND USER ACCEPT IT, SUBJECT TO THE FOREGOING CONDITIONS OF SALE AND WARRANTY.

TOLERANT ORNAMENTAL SPECIES:

This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

SCIENTIFIC NAME	COMMON NAME	SCIENTIFIC NAME	COMMON NAME	SCIENTIFIC NAME	COMMON NAME
Abies spp.	Fir species** (Balsam, Frasier, Noble, etc.)	Hedera helix	English Ivy	P. taeda	Loblolly Pine
Abelia grandiflora	Abelia: Sherwood	Helianthemum spp.	Sunrose	P. thunbergiana	Japanese Black Pine
Acer palmatum	Japanese Maple	Hemerocallis spp.	Daylily: Aztec Gold, Stella de Oro, Tender Love	P. virginiana	Virginia Pine
A. Platanoides	Norway Maple	Hibiscus	Rose of Sharon**	Pistachio spp.	Pistachio*
Achillea spp.	Yarrow: King Edward	Hibiscus spp.	Mallow: Disco Belle White	Pittosporum rhombifolium	Queensland Pittosporum
Actinidia chinensis	Kiwi*	Hibiscus rosa-sinensis	Chinese Hibiscus	P. tobira	Japanese Pittosporum
Agapanthus africanus	Lily-of-the-Nile (African Lily)	Hosta plantaginea	Hosta, Plantain Lily (Fragrant)	Podocarpus macrophyllus	Japanese Yew
Agapanthus orientalis		H. Sieboldiana	Hosta, "Seersucker"	Prunus laurocerasus	English Laurel
Akebia quintata	Five-leaf or Chocolate Vine		Houttuynia cordata var. variegata	Prunus spp.	Almond, Apricot, Nectarine, Peach, Plum and Prune*
Allium cernuum	Lady's Leek, Nodding Onion	Hydrangea macrophylla	Bigleaf Hydrangea	Pseudotsuga menziesii	Douglas Fir**
Anemone hybrida	Japanese Anemone	Ilex Bernyi	Holly	Pyracantha coccinea	Firethorn (Scarlet)
Aquilegia spp.	Aquilegia: Red and Gold	Ilex cornuta	Chinese Holly	P. fortuneana	Firethorn
Arctostaphylos densiflora	Vine Hill Manzanita	I. crenata	Japanese Holly	P. koidzumii	Firethorn
Arctotheca calendula	Cape Weed	I. opaca	American Holly	Pyrus spp.	Bradford Pear spp.
Artemisia spp.	Wormwood; Silver Mound, Castle	I. vomitoria	Yaupon Holly, Schillings	Quercus shumardii	Oak, Shumard's Red
Aster spp.	Aster: Bonnie Blue, Purple Dome		Inula ensifolia	Quercus spp.	Oak species
	Aster X Frikarti	Iris ensata	Sword-Leaved Iris; Jodlesong	Raphiolepis indica	Indian Hawthorne
Athrium filiz-femina	Lady Fern; Fern Lady	Iris siberica	Siberian Iris; Cabernet	Raphiolepis umbellata	Yedda Hawthorne
Aucuba japonica	Japanese Aucuba	Iris spp.	Iris species**	Rhododendron	'Coral Bells,' 'Formosa,' 'Hino-crimson,' 'PJM'
Begonia spp.	Fibrous Begonia: Hardy Grandis	Jasminium nudiflorum	Winter Jasmine	(including Azalea)	'Delaware Valley White,' 'Roseum Elegans,' 'Flame Creeper,' 'Girard Crimson,' 'Wakeiebisu'
Berberis gladywynensis	Barberry	Juniperus chinensis	Chinese Juniper; Nick's Compact, Parsonii	Rosa banksiae	Lady Bank's Rose
B. Julianae	Wintergreen Barberry	J. conferta	Shore Juniper; Blue Pacific	Rosmarinus officinalis	Rosemary*
B. mentorensis	Mentor Barberry	J. davurica	Parsoni	Rudbeckia spp.	Black-Eyed Susan; Goldsturm
B. Thunbergii	Japanese Barberry	J. horizontalis	Creeping Juniper	Santolina virens	Santolina
B. Verruculosa	Warty Barberry	Juglans spp.	Walnut*	Saxifraga spp.	Saxifrage; Purple Dome
	Bergenia cordifolia	Justicia brandegeana	Shrimp Plant	Scabiosa spp.	Pin cushion Flower
		Lagerstromia indica	Crape Myrtle	Sedum album	Stonecrop
Boltonia asteroides	Snowbank	Lagerstromia indica X Fauriei	Crape Myrtle; Tuscarora	Sedum caucicola	Stonecrop; Lidakense
Buddleia davidii	Butterfly-Bush (Dwarf Blue); Royal Red	Lantana montevidensis	Weeping Lantana	Sedum dasyphyllum	Stonecrop
Buxus microphylla	Japanese Boxwood	Lavender spp.	Lavender; Munstead	Sedum spuriatum	Stonecrop; Dragon's Blood
Callistemon viminalis	Weeping Bottlebrush	Leonopodium alpinum	Edelweiss	Spiraea bumalda	Spirea; Anthony Waterer
Calluna vulgaris	Scotch Heather	Ligustrum amurense	Amur Privet	Syzygium paniculatum	Australian Brushcherry
Campanula carpatica	Tussock Bellflower, (White Clips)	Ligustrum sinense	Chinese Privet; Variegata	Taxus cuspidata	Japanese Yew
Campis X Tagliabuana	Trumpet Flower, Madame Galen	L. japonicum	Japanese Privet	T. media	Yew
Carpobrotus edulis	Hottentot Fig (Ice Plant)	L. lucidum	Glossy Privet (Waxleaf)	Teucrium spp.	Germander
Cassia artemisoides	Feathery Cassia	Lilium spp.	Lily; Jazz	Thalictrum dipterocarpum	Meadow Rue
Ceanothus rigidus	Wild Lilac	Lirioppe muscari	Big Blue Lily, Silver Mound, Evergreen Giant	Thuja occidentalis	American Arborvitae
	Ceratostigma plumbagonoides	Lirioppe muscari var. variegata		Trachelospermum asiaticum	Star Jasmine
Chamaecyparis pisifera	False Cypress	Lirioppe spicata	Lirioppe, Creeping	Tsuga canadensis	Canada Hemlock
Cleyera japonica	Cleyera	Lonibelia Cardinalis	Cardinal Flower, Indian Pink	Tulipa spp.	Tulip species
Citrus spp.	Citrus species*	L. tatarica	Japanese Honeysuckle	Veronica spp.	Veronica, Speedwell; Sunny Border
Coreopsis spp.	Coreopsis (Calliopsis) Early Sunrise, Moonbeam	Lorapetalum chinense	Tatarian Honeysuckle	Viburnum japonicum	Japanese Viburnum
Cornus florida	Flowering Dogwood	Lythrum spp.	Burgundy	V. odoratissimum	Sweet Viburnum
C. Stoloniifera	American Dogwood	Magnolia spp.	Magnolia species**	V. plicatum	Japanese Snowball
Cortaderia selloana	Pampas Grass	Maleophora luteola	Ice Plant	V. rigidum	Canary Island Viburnum
Cotoneaster apiculatus	Cranberry Cotoneaster	Malus spp.	Crabapple*	V. suspensum	Arrowwood Viburnum
C. buxifolius	Cotoneaster	Miscanthus sinensis	Yaku Jima, Silberfeder	V. tinus	Laurustinus
C. dammeri	Bearberry Cotoneaster	Nandina domestica	Heavenly Bamboo	V. trilobium	Cranberry Bush
C. Microphyllus	Rockspray Cotoneaster	Narcissus spp.	Narcissus species**	V. wrightii	Leatherleaf Viburnum
C. reticulatus	Hawthorne	Nerium oleander	Oleander	Vinca major	Greater Periwinkle, Vinca
C. reticulatus	Lucifer	Olea europaea	Olive*	Vinca Minor	Common Periwinkle, Vinca
Cupressus sempervirens	Italian Cypress	Ophiopogon japonicus	Mondo Grass	Vitis spp.	Grape*
Delosperma alba	White Trailing Ice Plant	Osmanthus heterophyllus	Osmanthus (False Holly); Gulf Tide	Weigela florida	Old Fashioned Weigela
Delosperma spp.	Cooperi Pink	Osteospermum fruticosum	Trailing African Daisy	Yucca aloifolia	Spanish Bayonet, Yucca
Delphinium spp.	Larkspur: Blue Elf	Oxydendrum arboretum	Sourwood	Y. filametosia	Adam's Needle, Yucca
Dianthus deltooides	Dianthus: Maiden Pinks "Zing"	Paeonia suffruticosa	Tree Peony		
D. gratianopolitanus	Cheddar Pink	Pennisetum setaceum	Fountain Grass (Dwarf)		
Daonea viscosa	Hop Bush	Perovskia atriplicifolia			
Echinacea purpurea	Coneflower, Purple; Magnus	Persea americana	Avocado*		
Elaeagnus pungens	Silverberry	Photinia Frasier	Frasier's Photinia (Redtip)		
Euonymus fortunei	Wintercreeper	Physotegia virginiana	Dragonhead, False; Vivid		
E. japonica	Evergreen Euonymus	Picea spp.	Spruce species** (Colorado Blue, Norway, etc.)		
E. kiautschovica	Spreading Euonymus	Pieris japonica	Lily-of-the-Valley Bush		
Fatsia japonica	Japanese Aralia	Pinus brutia	Calabrian Pine		
Forsythia intermedia	Border Forsythia	P. canariensis	Canary Island Pine		
Forsythia suspensa	Weeping Forsythia	P. Elliotii	Slash Pine		
F. viridissima	Greenstem Forsythia	P. halepensis	Aleppo Pine		
Gaillardia spp.	Gaillardia, Blanket Flower: "Goblin"	P. nigra	Austrian Pine		
Gardenia jasminoides	Gardenia, Cape-Jasmine	P. palustris	Longleaf Pine		
	Gaura spp.	P. radiata	Monterey Pine		
Gentiana dahurica	Gentian	P. strobus	Eastern White Pine		
Geranium cinereum	Cranesbill	P. sylvestris	Scotch Pine		
Gladiolus spp.	Gladiolus species**				
Gypsophila repens	Baby's Breath				

*Do not use on food producing trees, vines or plants.

**Not for use on container grown plants.